



Quality of life in pre- and post-treatment among obstetric fistula patients at Kisii Hospital, Kenya

Judy Mwangi¹, Stephen Mutiso² and Ruchi Puri³, Jennifer Gatebi⁴

Abstract

Background Urinary incontinence has been noted to be a major barrier to social activities, entertainment or physical recreation. Ample evidence suggests that urinary incontinence affects a person's quality of life.

Objective To establish the impact of a combined surgical procedure and health promoting intervention to quality of life in women treated for obstetric fistula.

Method This prospective study focused on women undergoing treatment in Kisii Level Five Hospital in Nyanza Region, Kenya in September 2009. Treatment comprised surgical care, physiotherapy, health education and psychosocial support. The outcome parameters were self-reported incontinence and quality of life measured by the psychometric tool; King Health Questionnaire (KHQ), obtained upon admission and at six months follow-up. Ethical accept was obtained from the AMREF Scientific and Ethical Review Committee.

Results 53 women were interviewed at baseline and 40 were followed up after six months. 27 of 40 patients (68%) had vesicovaginal fistula, the other 12 patients (30%) had recto-vaginal fistula, while 1 (3%) patient had both. Full continence was achieved for 26 of the 40 (65%), and an additional 9 patients (23%) had improved from continuous to stress incontinence. Only 5 of 40 patients had failed repair ($p < 0.05$). The KHQ outcomes showed significant improvements for all dimensions of quality of life measure, except personal relationships.

Conclusion Women with obstetric fistula were found to drastically improve the continence and quality of life after undergoing a combined surgical and health promoting programme in hospital.

About the AUTHORS

1 African Medical and Research Foundation (AMREF), Nairobi, Kenya.

2 Kenyatta University, Nairobi, Kenya.

3 Department of Obstetrics and Gynaecology, Hubert-Young Center for Global Health, Duke Global Health Institute, Duke University Medical Center, USA

4 African Medical and Research Foundation (AMREF), Nairobi, Kenya

Contact:
Judy W Mwangi
juddys2@yahoo.com

Introduction

Health Related Quality of Life (HRQoL), a multidimensional construct referring to patients' perceptions of the impact of disease and treatment on their physical, psychological and social function and well-being (1), is crucial in the assessment of health care interventions (2). Urinary incontinence, in whichever form, sweepingly affects the life of the patients. Causes of urinary incontinence in women include childbirth, menopause and surgery (3). Vesico-vaginal fistula occurs mostly when labour is prolonged/obstructed forming an abnormal opening between a woman's bladder and vagina, and subsequent urinary incontinence (4). A similar pathophysiology exists for recto-vaginal fistula, which is followed by faecal incontinence. The two conditions may co-exist.

It impacts not only the physiological and physical, but also the psychological

realms of a person's life (5). It is conceived as a lack of health, which generates feelings of anger and sadness, as well as embarrassment. There is also the stigma associated with urinary/stool incontinence. Due to the smell of urine, patients avoid social gatherings and lose self-confidence, which has a proportional impact on their social interactions, their sexual life and emotional health (5;6). Obstetric fistula survivors who have been shunned and isolated typically experience intense feelings of shame, self-loathing and depression (8). Obstetric fistula survivors often avoid social contact because of feelings of shame, which negatively influences their quality of life (QoL) (3;9;10).

Up to the present, only a few studies have investigated QoL after fistula surgery so there is paucity of information. Overall, the existing studies show that the symptoms in most patients are reduced after treatment, which leads to an increase of



Research and Best Practice

different aspects of well-being (5).

The paper aims to establish the impact of a combined surgical and health promoting intervention to quality of life in women treated for obstetric fistula in Kisii Level Five Hospital, Kenya in 2009. It is anticipated that the study findings will be used to contribute towards policy advocacy and improvement in future programming of maternal and reproductive health initiatives in the country.

Methods

The study used a prospective before-and-after design. The outcome parameters were self-reported incontinence and QoL. A structured interviewer-administered tool was used to collect demographic data, while a King Health Questionnaire (KHQ) tool was utilised to gather QoL measures from all consenting women on admission and six months later during a follow-up visit at the hospital. The KHQ is a psychometric tool designed to assess the impact of urinary incontinence on QoL in women. The KHQ contains 21 questions categorised into nine domains (general health perception, incontinence impact, role limitations, physical limitations, social limitations, personal relationships, emotions, sleep/energy, and severity of urinary symptoms) (10). Weighted summary scores in each domain range from 0 to 100, with higher scores indicating greater impairment. The last part of the questionnaire was a list of bladder problems plus a category covering other options. These items are not summed to form a domain score. However, only the first eight of KHQ dimensions were utilised for this study, as women with fistula were incontinent throughout, thus leaving the last part of KHQ inapplicable.

Patients

At Kisii Level Five Hospital in Nyanza Region, Kenya, a total of 53 women underwent repair for obstetric vesico-vaginal and/or recto-vagina fistula in September, 2009. 40 of the patients (75%) were followed up after six months. The others were lost to follow up as they hailed from distant areas and were referred to other health facilities nearest to them for further management. 27 of 40 (68%) patients had vesico-vaginal fistula, 12 (30%) patients had recto-vaginal fistula, while 1 (3%) had both. No other medical conditions were noted. The median age of the women was 28 years (15-70); age at marriage 18 years (15-25), while age at birth of first child was 16 years (13-24). The number of living children ranged from 0-8 (1 in median), but 11 (28%) had no living children. Primary repair cases were 34 (85%), while the rest, 6 (15%), were repeated repair cases. Other patients' characteristics are outlined in Table 1.

Table 1 Characteristics of the 40 patients followed-up

| | Patients n=40 | 100 % |
|---|------------------|-------|
| Marital status | | |
| Married | 25 | 62.5 |
| Single | 9 | 22.5 |
| Divorced/Separated/widowed | 6 | 15.0 |
| Education | | |
| None | 5 | 12.5 |
| Primary | 27 | 67.5 |
| Secondary | 6 | 15.0 |
| Tertiary | 2 | 5.0 |
| Employment status | | |
| Working class | 2 | 5.0 |
| Farmer/business | 18 | 45.0 |
| Housewife | 20 | 50.0 |
| Social support following fistula | | |
| Partner | 6 | 15.0 |
| Close relatives | 24 | 60.0 |
| Distant relatives | 2 | 5.0 |
| None | 8 | 20.0 |
| Duration lived with fistula (years) | | |
| <5 | 12 | 30.0 |
| 5-10 | 15 | 37.5 |
| 10+ | 13 | 32.5 |
| Baby outcome | | |
| Alive | 11 | 27.5 |
| Stillbirth | 29 | 72.5 |
| Number of children following fistula | | |
| None | 33 | 82.5 |
| 1+ | 6 | 15.0 |
| Unknown | 1 | 2.5 |

Health promotion activities carried out

Apart from surgery, the women admitted also underwent medical follow up at two weeks and one month following repair. Physiotherapy was offered to improve urinary incontinence, while health education was carried out to inform and empower the women and their family. The health education also focused on informing on appropriate future maternal health care and hygiene practices to promote quick recovery and prevent recurrence of the condition. Psychotherapy was carried out in all women due to high levels of depression noted in women suffering from fistula and experienced trauma following difficult child birth, child loss, occurrence of fistula and subsequent marital breakdown and stigma (Table 2).



Research and Best Practice

Table 2 Combined surgical and health promotion activities

| Interventions (general) | Interventions (specific) | Time period |
|---------------------------|--|--|
| Surgery/medical follow up | Fistula repair | Within the first week of screening/admission |
| | Catheterisation | At surgery |
| | Catheter removal | 2-weeks following surgery |
| | Medical review | Within one month following surgery |
| Physiotherapy | Pelvic floor exercises | Before and after surgery |
| Health education | Information on cause and prevention of obstetric fistula | Before and after surgery |
| | Family planning | |
| | Future pregnancies | |
| | Antenatal care and delivery | |
| | Sexual practices | |
| Psychotherapy | Group counseling | Before and after surgery |
| | Couple counseling | |
| | Individual counseling | |

Ethics

The patients participated after informed consent. Ethical review was obtained from the African Medical and Research Foundation (AMREF) Scientific and Ethical Review Committee.

Statistics

As the variables are not normally distributed, non-parametric statistical analyses have been used (Fisher's exact test for comparison of frequencies, and Wilcoxon test for paired data). A p-value < 0.05 was considered significant. The material seemed too small for sub-group analyses. Identification of patient characteristics associated with specific outcomes was carried out by univariate analyses, followed by a multivariate analysis. SPSS version 17.0 software was used for the analyses.

Results

Prior to intervention, 100% of the women were continuously incontinent. Significant improvement was seen after the combined intervention. Full continence was achieved for 26/40 (65%), and additional 9 patients (23%) had improved from continuous to stress incontinence. Only 5 of 40 patients had failed repair ($p < 0.05$).

Univariate analyses revealed that foetal outcome was

associated with incontinence, as those who delivered stillborn babies were more likely to report incontinence ($p=0.043$). None of the women with liveborn babies had self reported incontinence. No other associations between patient characteristics and outcomes were found, thus a multivariate analysis was not carried out. The KHQ outcomes at six months post-repair showed significant improvements in all the dimensions, except from personal relationships. The details are presented in Table 3.

Table 3 Kings Health Questionnaire (KHQ) scores

| Dimension | Median Score Pre-surgery (range) | Median Score Post-surgery (range) | P-value (Wilcoxon rank test) |
|----------------------------|----------------------------------|-----------------------------------|------------------------------|
| General health perceptions | 50 (0-100) | 25 (0-100) | 0.000 |
| Incontinence impact | 67 (0-100) | 33 (0-100) | 0.000 |
| Role limitations | 83 (0-100) | 0 (0-100) | 0.000 |
| Physical limitations | 100 (0-100) | 0 (0-83) | 0.000 |
| Social limitations | 67 (0-100) | 22 (0-100) | 0.000 |
| Personal relationships | 50 (0-100) | 67 (0-100) | 0.193 |
| Emotions | 100 (11-100) | 28 (0-100) | 0.000 |
| Sleep/energy | 67 (0-100) | 16 (0-100) | 0.000 |

Discussion

This study showed that almost nine of ten patients with obstetric fistula had effect of the combined surgical and health promoting programme on their incontinence and about two of three patients became fully continent. Furthermore, QoL increased dramatically for seven out of eight of the dimensions compared to baseline before intervention.

The patients in this study had background characteristics typical of fistula patients documented in Kenya and other parts in developing countries (11-15). The African Medical and Research Foundation (AMREF) is an African-based Non-Governmental Organization that has been implementing the Safe Motherhood Initiative including prevention and treatment of obstetric fistula both at community and health facility level since 2006. Prior to this project, the Safe Motherhood Initiative had been running obstetric fistula outreaches and camps since 1995 in the remote health facilities, where healthcare is almost inaccessible to the women due to poor road infrastructure and long distances to these facilities. However, that activity included a surgical approach, exclusively, with no follow-up or non-surgical interventions.



Research and Best Practice

In the present study, improvement in QoL following intervention was recorded in the emotion dimension. This dimension focused on three sub-components; feelings of depression, anxiety/nervousness and feeling bad about one-self. In general, depression is characterized by feelings of sadness, emptiness, dissatisfaction, lowered self-esteem, inactivity, self-depreciation and despair (16). A previous study has shown that most women suffering from fistula also suffer from depression (11). Unless the women were aware that fistula could be treated and gained access to provide subsidised treatment and care, women would live with the fistula for life (11). Lack of knowledge about the root cause of fistula persists in many communities, heightens the level of stigma and discrimination, in addition to increasing delays in seeking surgical repair (14). These factors were reflected in the women interviewed in the present study. About two thirds of the women reported to have lived with fistula for over 10 years, with the longest duration recorded in a patient in this study being 40 years.

In African culture, woman's worth can easily be determined by the number of children that she bears for her husband and therefore a barren woman is seen as a disgrace and of no use to her husband (18), which could aggravate feelings of anxiety and depression. This reflects the scenario in the study group, in which the mothers who suffered a traumatic delivery, mostly lost their children at birth, had no other living children and also had not been able to get more children following fistula; factors that could have easily contributed to exacerbate stigma, thus heightened depression.

Loss of continence and inability to maintain hygiene standards among these women may also be accompanied by alienation from family and friends and this may be additionally detrimental to the patient's self-confidence (19-20). Interestingly, personal relationships were not improved by the combined surgical and health promoting programme. Both social limitations and personal relationships had the lowest score at follow-up. These two dimensions involve other persons, and the women may require some more time to regain the level of activity they engaged in prior to the occurrence of fistula. There was no significant association between the social demographic characteristics and incontinence. However, the foetal outcome was associated with incontinence; those who delivered stillborn babies were more likely to report incontinence. Foetal outcome is implicated with prolonged/obstructed labour, which denotes a higher chance of developing fistula/incontinence.

This study was not sizable for sub-analysis. Researchers from Ethiopia found no association between social

demographics and depression markers, but noted differences in relation to continence (21). In Kenya, the present authors observed no significant difference in depression between primary and repeated cases (14).

The strength of the study was its real life setting and that it did not exclude important patient groups on forehand. The relatively high follow-up rate is also considered a strength.

This study has the following limitations; 1) there was no control group; 2) the sample size was very small, which could lead to type-2 failures; 3) the patients had no follow-up performed later than the six months follow-up. In addition, the significance found in this study could be due to a type-1 failure, which could be reduced by repeating the study. The measurement of personal relationships focused on relationships with a partner, sex life and family life, and the answers may be influenced by the advice given to abstain from sex for six months following surgery. This advice guarantees time for the fistula to heal, but also affects the sexual relationships with the partners further. The bias and limitations call for careful interpretation and caution in generalisation of the results. On the other hand, until larger studies in higher quality design may indicate otherwise, implementation of the combined programme under further monitoring seems indicated, because obstetric fistula is very important to treat due to its impact on the QoL of the patient. From this study, it is not possible to identify which or if any specific elements in the combined surgical and health promotion programme are of most importance. This is, however, a general problem for all combined interventions, and therefore the attention should be on implementing the complete programme. When monitoring the implementation, more data will be gathered to further evaluate the effect and to perform relevant sub-group analyses.

Conclusion

In conclusion, women with obstetric fistula were found to drastically improve their continence and quality of life after undergoing a combined surgical and health promoting programme in hospital. We recommend early repair of fistula to prevent prolonged diminishing of quality of life in the patients. In addition, to promote prevention of development of obstetric fistula, we also recommend health promotion programmes at community and health facility level for the benefit of the women, the families and the society as a whole.



Research and Best Practice

Acknowledgments

African Medical and Research Foundation (AMREF) and Kisii Level Five Hospital.

Contribution Details

JM and SM designed the study, SM, JM and RP performed the research, JM and JG collected the data, JM and JG analyzed the data, JM and SM wrote the paper, and SM, RP & JM edited the paper.

Competing Interests: None declared

References

- (1) Rowles SV, Prieto L, Badia X, Shalet SM, Webb SM, Trainer PJ. Quality of life (QOL) in patients with acromegaly is severely impaired: use of a novel measure of QOL: acromegaly quality of life questionnaire. *J Clin Endocrinol Metab* 2005; 90:3337-41.
- (2) Feeny D, Furlong W, Saigal S, Sun J. Comparing directly measured standard gamble scores to HUI2 and HUI3 utility scores: group- and individual-level comparisons. *Soc Sci Med* 2004; 58:799-809
- (3) Broome BAS. The impact of urinary incontinence on self-efficacy and quality of life. *Health and qual of life outcomes* 2003; 1:35.
- (4) Wall LL. Obstetric vesicovaginal fistula as an international public-health problem. *Lancet* 2006; 368:1201-09.
- (5) Swithinbank L, Abrams P. The impact of urinary incontinence on the quality of life of women. *World J Urol* 1999; 17:225-9.
- (6) Lalos O. Impact of urinary and climacteric symptoms on social and sexual life after surgical treatment of stress urinary incontinence in women: a long-term outcome. *J Advances Nursing* 2001; 33:316-27.
- (7) UNFPA and Engender Health: Obstetric Fistula Needs Assessment Report 2003; ISBN: 0-89714-661-1.
- (8) Ouslander J, Schnelle J. Incontinence in the nursing home. *Ann Intern Med*

1995; 122:438-49.

- (9) Hunskar S, Vinsnes A. The quality of life in women with urinary incontinence as measured by the sickness impact profile. *J Am Geriatr Soc* 1992; 39:378-82.
- (10) Kelleher CJ, Cardozo LD, Khullar V, Salvatore S: A New Questionnaire to Assess the Quality of Life of Urinary Incontinent Women. *Br J Obstet Gynaecol* 1997; 104:1374-9.
- (11) UNFPA and Kenya Ministry of Health: Needs Assessment of obstetric fistula in selected districts of Kenya. Nairobi, 2004. <http://www.fistulanetwork.org>. Accessed Sep 20, 2010.
- (12) Muleta M. Obstetric Fistula in Developing Countries: A Review Article. *J Obstet Gynaecol Can* 2006; 28:962-6.
- (13) Raassen TJ, Verdaasdonk EG, Vierhout ME. Prospective results after first-time surgery for obstetric fistulas in East African women. *Int Urogynecol J Pelvic Floor Dysfunct*. 2008; 19:73-9.
- (14) Weston K, Mutiso S, Mwangi JW, Qureshi Z, Beard J, Venkat P. Depression among women with obstetric fistula in Kenya. *Int J Gynaecol Obstet* 2011; 115:31-3.
- (15) Wall LL, Arrowsmith SD, Briggs ND, Browning A, Lasey A. The obstetric vesicovaginal fistula in the developing world. *Obstet Gynecol Surv* 2005; 60 (7 Suppl 1):S3-S51.
- (16) Lewis S, Grainger RD. Manual of psychosocial nursing interventions: Promoting mental health in medical-surgical settings. In: *The patient with depression*. Philadelphia, W.B. Saunders Co. 1989.
- (17) Bandura A. Self-efficacy mechanisms in human agency. *Am Psychol* 1982; 37:122-47.
- (18) Bacon, C. (Ed.) 2003. *Obstetric Fistula Needs Assessment Report: Findings from Nine African Countries*. NYC, USA: UNFPA and Engender Health.
- (19) Fultz N, Herzog R. Self-Reported Social and Emotional Impact of Urinary Incontinence. *J Am Geriatr Soc*. 2001; 49:892-9.
- (20) Wall LL, Arrowsmith S, Briggs NS, Lasey A, 2001. Urinary incontinence in the developing world: The obstetric fistula. *Proceedings of the 2nd International Consultation on Urinary Incontinence*, Jul. 1-3, Paris, France, 1-40.
- (21) Browning A, Fentahun W, Goh J. The impact of surgical treatment on the mental health of women with obstetric fistula. *BJOG: An Int J of Obste & Gyneco* 2007; 114:1439-41.

Publish in Clinical Health Promotion

Clinical Health Promotion is always looking for new interesting research for the Journal

We want your research to have the highest impact and the widest audience possible, which is why both the printed and online version is free of charge.

Publishing in Clinical Health Promotion is free of charge and all contributions should be submitted via www.clinhp.org

For more information about Clinical Health Promotion and for details concerning submission please visit www.clinhp.org.

